

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : <b>A61K 7/28</b>	<b>A1</b>	(11) International Publication Number: <b>WO 97/06775</b>
		(43) International Publication Date: 27 February 1997 (27.02.97)

(21) International Application Number: PCT/DK96/00350

(22) International Filing Date: 19 August 1996 (19.08.96)

## (30) Priority Data:

0926/95	18 August 1995 (18.08.95)	DK
1048/95	20 September 1995 (20.09.95)	DK

(71) Applicant (for all designated States except US): NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsvaerd (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): AASLYNG, Dorrit [DK/DK]; Novo Nordisk a/s, Novo Allé, DK-2880 Bagsvaerd (DK). TSUCHIYA, Rie [JP/DK]; Novo Nordisk a/s, Novo Allé, DK-2880 Bagsvaerd (DK).

(74) Common Representative: NOVO NORDISK A/S; Corporate Patents, Novo Allé, DK-2880 Bagsvaerd (DK).

(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

**Published***With international search report.*

(54) Title: TOOTH BLEACHING

## (57) Abstract

The object of the present invention is to provide safe products for bleaching teeth. Oral compositions comprising at least one oxidoreductase fulfil said criteria without the presence of peroxide or with only very little peroxide present. The invention also relates to an oral care product comprising the oral composition of the invention, a method of bleaching teeth, a method for using said oral care products and the use of oxidoreductases for oxidation of teeth stains.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**Title:** Tooth Bleaching

**5 FIELD OF THE INVENTION**

The present invention relates to an oral composition for bleaching teeth without the presence of peroxide or comprising only very little peroxide, an oral care product comprising the oral composition of the invention, a method of bleaching  
10 teeth, a method for using said oral care product and the use of oxidoreductases for oxidation of teeth stains.

**BACKGROUND OF THE INVENTION**

15 Most people want to have a mouth full of dazzling white teeth, as the teeth constitute an important part of the overall picture of the human face especially when smiling. In contrast hereto strongly discoloured teeth entail an unhealthy and in certain cases even repellent "look".

20 For many years, crowns or dentures were seen as the only means for avoiding e.g. the yellowing of teeth coming with age, discolouration of teeth due to long term intake of the antibiotic tetracycline during childhood, or the yellowish brown to black  
25 discolouration of teeth as a consequence of coffee drinking, tobacco smoking/chewing etc.

Even though techniques for bleaching teeth have been known for many years, tooth bleaching has never been used extensively,  
30 until recently.

The first published tooth bleaching technique used oxalic acid as the bleaching agent and was reported by Chaple in the 1877. Soon after, the first report of peroxide used as a bleaching  
35 agent was published by Harlan in 1884. Even though quite a number of bleaching agents have been suggested since then peroxide is today still the most commonly used agent.

### The structure of teeth

A mammal tooth is chiefly composed of four main constituents namely the "cementum", the "dental pulp", the "dentin", and the "enamel".

5

The "cementum" is bonelike tissue forming the outer surface of the root of the tooth. The "dental pulp" consists of sensitive tissue containing arteries, veins, and lymphatic and nerve tissue. The "dentin" which surrounds the dental pulp forms the major part of the tooth. The dentin is dense bonelike calcareous tissue. On the outside of the tooth there is a hard porous layer composed of hydroxyapatite mineral crystals having a natural opaque white or off-white colour. This outer porous layer is called the "enamel".

15

### Tooth discolouration

Tooth discolouration can be caused by a variety of intrinsic and/or extrinsic influences. In general stains are divided into two main categories:

20 1) extrinsic stains and 2) intrinsic strains.

#### 1) Extrinsic stains

Extrinsic stains are mainly caused by the daily intake of substances, such as foods and beverages through the mouth, and/or the use of tobacco products etc. These substances tend to adhere to the enamel's hydroxyapatite structure and hereby discolour the teeth and/or reduce the whiteness of the teeth.

Over a period of years extrinsic stains may penetrate the enamel layer and gradually give intrinsic discolourations.

#### 2) Intrinsic stains

Intrinsic stains are the term used for stains which have penetrated the tooth structure (i.e. discolouration within the tooth matrix). Such stains can arise as described above or be caused by agents including haematological agents and certain drugs, or be due to dental pulp necrosis or developmental abnormalities.

35

For instance, degradation products from the body may cause discolorations. Excessive intake of tetracycline or fluoride during a long period of illness has been found to release degradation products into the dentinal tubules during the development of tooth enamel causing some degree of discolouration. The severity of such discolouration depends on the time and duration of intake of the medicine.

10 Further, dental pulp necrosis entails Haemorrhagic discolorations and is a result of blood degradation. If the pulp necrosis is caused by e.g. caries degradation of proteins the tooth/teeth will become greyish-brown.

15 In the case of traumatic pulp death the tooth will turn yellow-brown. It is believed that such a discolouration is caused by haemolysis of red blood cells entering the dentinal tubules.

20 Tooth bleaching

Before conducting tooth bleaching it is important to assess the type of stain, as different types of stains need different approaches and/or bleaching agents.

25 Certain extrinsic stains, which occur on the surface or sub-surface of the teeth, can be removed by regular intense mechanical brushing of the teeth with cleansing agents containing abrasives and surfactants. However, not all extrinsic stains can be removed this way and require bleaching agents which inhibit non-enzymatic browning reactions.

Intrinsic stains are located in the tooth matrix and cannot be removed or prevented by intense mechanical brushing of the teeth. Removal of such discolourations requires bleaching agents capable of penetrating into the teeth structure. Hydrogen peroxide is an example of such an agent, which can be used for both extrinsic and intrinsic stains.

Hydrogen peroxide can be used for many types of stains e.g. stains residing in the dentin, such as stains caused by tetracycline.

- 5 Even though hydrochloric acid is not regarded as being a bleaching agent it is known to be capable of removing stains caused by fluorosis, as it dissolves the surface of the teeth.

10 However, the use of such agents can inflict severe tooth damage or at least irritation in the oral cavity. Consequently, such agents are for safety reasons not suitable for "home-use" by the private consumer and should only be used precautiously by professionals.

15 Bleaching techniques

Bleaching techniques are usually divided into two main categories:

- a) non-vital bleaching techniques, and
- b) vital bleaching techniques.

20

a) Non-vital bleaching

The non-vital techniques give the most effective results but also have the greatest potential hazard. One non-vital bleaching technique uses sodium perborate and 35% hydrogen peroxide as the active ingredient.

25

b) Vital bleaching

Products sold for vital bleaching techniques can be divided into three main groups a) "in-office" bleaching products, b) dentist prescribed, home applied bleaching products, and c) over-the-counter bleaching kits.

30

For further information concerning the categorisation of tooth bleaching products and techniques we refer to Van B. Haywood, (1992), Periodontology and Restorative Dentistry, p. 142-149.

35

One of the most commonly used "in-office"-techniques combines the use 30% hydrogen peroxide with heat and light treatment to

speed up the oxidation reaction (i.e. the removal of stains).

Another method, using a "dentist prescribed, home-applied"-bleaching product, involves the use of 10% urea peroxide (carbamide peroxide). The teeth are bleached in a mouth tray, containing the bleaching agent, placed upon the teeth of the patient.

Over-the-counter kits which can be used for bleaching teeth include products such as toothpastes and mouth washes having from 3% to 6% hydrogen peroxide and are sold directly to the consumer, without prescription by a dentist.

#### Comments to prior art

Most of the above mentioned prior art methods/techniques involve the use of peroxides and/or other agents in concentrations which are not safe for "home-use" by the private consumer due to the risk of damaging the teeth and/or the oral tissue.

Furthermore, effective concentrations of e.g. hydrogen peroxide exceed the allowed limits in certain countries.

Products comprising low concentration of bleaching agents, such as hydrogen peroxide, are considered to have slow bleaching effect.

Therefore, there is a need for providing safe tooth bleaching compositions, which do not comprise harmful concentrations of peroxide and/or other hazardous agents. It is further desirable that such tooth bleaching compositions can be used as components in conventional oral care products for "home-use" by the private consumer.

**SUMMARY OF THE INVENTION**

The object of the present invention is to provide safe tooth bleaching products, which do only comprise very low concentrations of peroxides or/and other hazardous bleaching agents.

The present inventors have surprisingly found that tooth bleaching compositions comprising at least one oxidoreductase have a good bleaching effect without causing the problems that high concentrations of peroxides and other agents may cause.

Consequently, in the first aspect the object of the present invention is to provide an oral composition for bleaching teeth comprising at least one oxidoreductase, such as a laccase or a related enzyme, and/or an oxidase and/or a peroxidase.

In the second aspect the invention relates to an oral care product, comprising an oral composition for bleaching teeth of the invention.

The invention also relates to a method for bleaching discoloured or stained teeth, and a method for using an oral composition or an oral care product in the oral cavity.

Finally the invention relates to the use of oxidoreductases for vital and non-vital bleaching of teeth.

30

**DETAILED DESCRIPTION OF THE INVENTION**

The object of the present invention is to provide safe tooth bleaching products which do only comprise very low concentrations of peroxides or/and other hazardous agents.

In an embodiment of the invention it is even possible to omit peroxide and/or other equivalent agents completely from the



composition.

In the context of the present invention the term "bleaching" of teeth is equivalent with the terms "whitening" or "brightening" of teeth.

The term "low concentrations" of e.g. peroxide means concentration of from 0% (no peroxide present) to about 1% peroxide, calculated on the basis of the weight of final oral composition or oral care product.

All concentrations mentioned in the connection with the present invention are calculated in weight per cent.

Safe oral compositions and oral care products of the invention may comprise from 0% to 0.5% peroxide, such as less than about 0.3%, which may be about 0.1% peroxide. Preferably the concentration of peroxide lies from 0% to 1%.

It is to be understood that even though less than 1% of peroxide is needed for obtaining a tooth bleaching effect of the invention, it is contemplated to add concentration of peroxide that is higher than 1%, such as 3% or 6% and even 10% calculated on the basis of the weight of the final oral composition or oral care product.

However, if peroxide is present in higher concentrations it may damage the oral tissue. Further, the stain oxidising enzyme may be inactivated at such concentrations.

The use of oral care product comprising an oral composition of the invention facilitates the bleaching of teeth. For instance oral care products of the invention in the form of a toothpaste or a mouth wash make it possible to incorporate tooth bleaching as a natural part of the daily teeth cleaning and/or mouth rinsing at home.

The present inventors have surprisingly found that safe tooth

bleaching compositions can be provided by adding an enzyme or enzyme system capable of oxidising the teeth stains.

5 In the first aspect the invention relates to an oral composition for bleaching teeth comprising at least one enzyme within the group of oxidoreductases as the active bleaching ingredient(s).

10 Oxidoreductases (i.e. enzymes classified under the Enzyme Classification number E.C. 1 (Oxidoreductases) in accordance with the Recommendations (1992) of the International Union of Biochemistry and Molecular Biology (IUBMB)) which are enzymes catalysing oxidoreductions.

15 Within the group of oxidoreductases enzymes are preferred which catalyse the oxidation of a substrate (an electron or hydrogen donor) by acting on oxygen ( $O_2$ ) and/or a peroxide as the acceptor. Such enzymes include enzymes classified within the enzyme classes comprising oxidases including E.C. 1.1.3.  
20 E.C. 1.2.3, E.C. 1.3.3, E.C. 1.4.3, E.C. 1.5.3, E.C. 1.7.3, E.C. 1.8.3, E.C. 1.9.3, laccases and related enzymes comprised in E.C. 1.10.3 and peroxidases in E.C. 1.11.

25 In the case of an enzyme acting on oxygen ( $O_2$ ) as the acceptor, said oxygen may be molecular oxygen supplied by the air.

30 If a peroxide, for instance  $H_2O_2$  or  $H_2O_2$ -generation compounds such as perborat and percarbonate, is added to the oral composition or oral care product it will be added in the above mentioned concentrations.

35 Enzymes' or enzyme systems' ability to fulfil the above criteria of oxidising teeth stains can be assayed by using the approach described below in the "Materials and Methods"-section.

According to the invention three types of oxidoreductases are

specifically contemplated:

a) Laccases or related enzymes, which act on molecular oxygen and yield water ( $H_2O$ ) without any need for peroxide (e.g.  $H_2O_2$ ),

b) Oxidases, which acts on molecular oxygen ( $O_2$ ) and yield peroxide ( $H_2O_2$ ), and

c) Peroxidases, which act on peroxide (e.g.  $H_2O_2$ ) and yield water ( $H_2O$ ).

Also enzyme systems which comprise a combination of the three types of enzymes are contemplated according to the invention. The enzyme systems may e.g. consist of a laccase and an oxidase; a laccase and a peroxidase; a laccase and an oxidase and a peroxidase; or an oxidase and a peroxidase.

Preferred are the below mentioned enzymes, especially recombinant and/or substantially purified enzymes.

In the context of this invention "laccases and related enzymes" include enzymes comprised by the enzyme classification E.C. 1.10.3.2 (laccases) and catechol oxidase enzymes comprised by E.C. 1.10.3.1, bilirubin oxidase enzymes comprised by the enzyme classification E.C. 1.3.3.5 and monophenol mono-oxygenase enzymes comprised by the enzyme classification E.C. 1.14.99.1.

Preferably, the laccase employed is derived from a strain of *Polyporus* sp., in particular a strain of *Polyporus pinsitus* or *Polyporus versicolor*, or a strain of *Myceliophthora* sp., e.g. *M. thermophila* or a strain of *Rhizoctonia* sp., in particular a strain of *Rhizoctonia praticola* or *Rhizoctonia solani*, or a strain of a *Rhus* sp., in particular *Rhus vernicifera*.

In specific embodiments of the invention the oxidoreductase is a laccase, such as a *Polyporus* sp. laccase especially the *Polyporus pinsitus* laccase (also called *Trametes villosa*

laccase) described in WO 96/00290 (from Novo Nordisk Biotec, inc.) or a *Myceliophthora* sp. laccase especially the *Myceliophthora thermophila* laccase described in WO 95/33836 (from Novo Nordisk Biotech inc).

5

Further, the laccase may be a *Scytalidium* sp. laccase, such as the *S. thermophilum* laccase described in WO 95/33837 (from Novo Nordisk Biotech inc.) or a *Pyricularia* sp. laccase, such as the *Pyricularia oryzae* laccase which can be purchased from  
10 SIGMA under the trade name SIGMA no. L5510, or a *Coprinus* sp. laccase, such as a *C. cinereus* laccase, especially a *C. cinereus* IFO 30116 laccase, or a *Rhizoctonia* sp. laccase, such as a *Rh. solani* laccase, especially the neutral *Rh. solani* laccase described WO 95/07988 (from Novo Nordisk A/S) having a  
15 pH optimum in the range from 6.0 to 8.5.

The laccase may also be derived from a fungi such as *Collybia*, *Fomes*, *Lentinus*, *Pleurotus*, *Aspergillus*, *Neurospora*, *Podospora*, *Phlebia*, e.g. *P. radiata* (WO 92/01046), *Coriolus* sp., e.g.  
20 *C. hirsitus* (JP 2-238885), or *Botrytis*.

Bilirubin oxidase may preferably be derived from a strain of *Myrothecium* sp., such as *M. verrucaria*.

25 Within the group of peroxidases classified under the Enzyme Classification number E.C. 1.11 (peroxidases) peroxidases (1.11.1.7) are especially contemplated.

The peroxidase of the invention may be derived from plants  
30 (e.g. horseradish peroxidase) or micro-organisms including fungi and bacteria such as a strain of *Coprinus* sp., such as *Coprinus cinereus* or *Coprinus macrorhizus*, or bacteria such as *Bacillus*, such as *Bacillus pumilus*.

35 In another embodiment of the invention the oxidoreductase is a peroxidase, such as a *Coprinus* sp. peroxidase, such as the *C. cinereus* peroxidase, such as *C. cinereus* IFO8371 or variants thereof described in WO 95/10602 (from Novo Nordisk A/S) being

very stable in the presence of hydrogen peroxide, or a haloperoxidase originating from a strain of *Curvularia* sp. such as a *C. verruculosa* haloperoxidase, in particular *C. verruculosa* CBS 147.63.

5

Oxidases which are contemplated include glucose oxidase (E.C. 1.1.3.4), hexose oxidase (E.C. 1.1.3.5), L-amino-acid oxidase (E.C. 1.4.3.2), xylitol oxidase, galactose oxidase (E.C. 1.1.3.9), pyranose oxidase (E.C. 1.1.3.10), alcohol oxidase  
10 (E.C. 1.1.3.13).

If a L-amino acid oxidase is used it may be derived from a *Trichoderma* sp. such as *Trichoderma harzianum*, such as the L-amino acid oxidase described in WO 94/25574 (from Novo Nordisk  
15 A/S), or *Trichoderma viride*.

A suitable glucose oxidase may originate from *Aspergillus* sp., such as a strain of *Aspergillus niger*, or from a strain of *Cladosporium* sp. in particular *Cladosporium oxysporum*,  
20 especially *Cl. oxysporum* CBS 163 described in WO 95/29996 (from Novo Nordisk A/S).

Hexose oxidases from the red sea-weed *Chondrus crispus* (commonly known as Irish moss) (Sullivan and Ikawa, (1973),  
25 Biochim. Biophys. Acts, 309, p. 11-22; Ikawa, (1982), Meth. in Enzymol. 89, carbohydrate metabolism part D, 145-149) oxidises a broad spectrum of carbohydrates, such as D-glucose, D-galactose, maltose, cellobiose, lactose, D-glucose 6-phosphate, D-mannose, 2-deoxy-D-glucose, 2-deoxy-D-galactose,  
30 D-fucose, D-glucuronic acid, and D-xylose.

Also the red sea-weed *Iridophycus flaccidum* produces easily extractable hexose oxidases, which oxidise several different mono- and disaccharides (Bean and Hassid, (1956), J. Biol.  
35 Chem, 218, p. 425; Rand et al. (1972, J. of Food Science 37, p. 698-710).

The broad substrate spectrum of hexose oxidase is advantageous

in the connection with tooth bleaching as the total amount of usable substrate (i.e. carbohydrate) present in the mouth is significantly greater than for related enzymes having more specific catalytic properties.

5

Another relevant group of tooth bleaching enzymes is xylitol oxidases (see e.g. JP 80892242) which oxidises xylitol, D-sorbitol, D-galactitol, D-mannitol and D-arabinitol in the presence of oxygen. A xylitol oxidase can be obtained from  
10 strains of *Streptomyces* sp. (e.g. *Streptomyces* IKD472, FERM P-14339) having a pH optimum at 7.5, is stable at pH 5.5 to 10.5 and at temperatures up to 65°C; properties very well suited for oral care compositions and products. Further, the  
15 substrate xylitol is not cariogenic (i.e. is not degraded in the mouth to compounds responsible for dental holes).

It can be advantageous to use enzyme(s) which can act on substrates which are not cariogenic (i.e. substrates which are not or is not immediately degraded into cariogenic substrates  
20 such as sucrose, glucose, fructose, maltose etc.).

Examples of such a substrate include amino acids, alcohol, sugar alcohol, such as xylitol, sorbitol etc.

25 Consequently, in a preferred embodiment the oral care composition comprises one or more of the above enzymes and a substrate which is not cariogenic.

It is also advantageous to use enzymes being substantially  
30 active at pHs prevailing in the mouth, i.e. between pH 5.0 to 9.0, preferably between pH 6.0 to 8.5, especially between pH 6.4 to 7.5.

The term "substantially active" enzyme means in this context  
35 that the enzyme(s) has(have) an relative activity (pH-optimum defines 100% at the same conditions) higher than 30%, better 50%, even better more than 70%, such as 80%, and in the best case up to about 100% of the activity at the pH optimum.

In the case of using laccase and a peroxidase the bleaching effect can be obtained by direct oxidation of the teeth stains or via a mediator.

5

Oxidases generate peroxide ( $H_2O_2$ ) *in situ* (i.e. in the oral cavity).

10

When using laccase and/or peroxidase, or an oxidase and a peroxidase no peroxide need to be present, while low concentrations of peroxide need to be added in the case of a peroxidase alone. However, it is also contemplated according to the invention to add low concentrations of peroxide when using laccase or oxidase.

15

It is to be understood that an advantage of the invention is that peroxide will not at any time be present in the oral cavity in concentrations which can inflict damage.

20

Compositions and products of the invention can be used safely without any major precautions by both the private consumer and professionals (i.e. dentists or the like) due to the low concentration or absence of hydrogen peroxide.

25

In a preferred embodiment the oral composition or oral care product of the invention comprises a redox mediator (sometimes called an enhancing agent or accelerator) which is an agent capable of enhancing the activity of oxidoreductases contemplated according to the invention.

30

The mediator may be any known mediator. Examples of such mediators are the following compounds: 2,2'-azino-bis(3-ethyl-benzothiazoline-6-sulfonate (ABTS); 6-hydroxy-2-naphtoic acid; 7-methoxy-2-naphtol; 7-amino-2-naphthalene sulfonic acid; 5-amino-2-naphthalene sulfonic acid; 1,5-diaminonaphthalene; 7-hydroxy-1,2-naphthimidazole; 10-methylphenothiazine; 10-phenothiazine-propionic acid (PPT); N-hydroxysuccinimide-10-phenothiazine-propionate; benzidine; 3,3'-dimethylbenzidine; 3,3'-

35

dimethoxybenzidine; 3,3',5,5'-tetramethylbenzidine; 4'-hydroxy-4-biphenylcarboxylic acid; 4-amino-4'-methoxystilbene; 4,4'-diaminostilbene-2,2'-disulfonic acid; 4,4'-diaminodiphenylamine; 2,7-diaminofluorene; 4,4'-dihydroxy-biphenylene; triphenylamine; 10-ethyl-4-phenothiazinecarboxylic acid; 10-ethylphenothiazine; 10-propylphenothiazine; 10-isopropylphenothiazine; methyl-10-phenothiazinepropionate; 10-phenylphenothiazine; 10-allylphenothiazine; 10-phenoxazinepropionic acid (POP); 10-(3-(4-methyl-1-piperazinyl)propyl)phenothiazine; 10-(2-pyrrolidinoethyl)phenothiazine; 10-methylphenoxazine; iminostilbene; 2-(p-aminophenyl)-6-methylbenzothiazole-7-sulfonic acid; N-benzylidene-4-biphenylamine; 5-amino-2-naphthalenesulfonic acid; 7-methoxy-2-naphthol; 4,4'-dihydroxybenzophenone; N-(4-(dimethylamino)benzylidene)-p-anisidine; 3-methyl-2-benzothiazolinone(4-(dimethylamino)benzylidene)hydrazone; 2-acethyl-10-methylphenothiazine; 10-(2-hydroxyethyl)phenothiazine; 10-(2-hydroxyethyl)phenoxazine; 10-(3-hydroxypropyl)phenothiazine; 4,4'-dimethoxy-N-methyl-diphenylamine; vanillin azine.

The amount of oxidoreductase(s) needed in an oral composition of the invention to obtain tooth bleaching depends on the particular compound employed, but ranges generally from 0.0001% to 20%, preferably from about 0.001% to about 10%, and most preferably from about 0.01% to about 5% by weight of the final composition.

In the preparation of an oral composition, the oxidoreductase(s) may for safety reasons be added as an essentially purified enzyme preparation. However, less purified oxidoreductases preparations can be used seen from a technical point of view.

The oral composition may be incorporated in products used for vital and/or non-vital tooth bleaching techniques. Examples of products suitable for vital tooth bleaching, having the main purpose of bleaching teeth, include products usually used by trained professionals in-office tooth bleaching and/or in the



so-called "dental prescribed, home-applied"-products.

Especially contemplated according to the invention are "over-the-counter"-products, which include tooth bleaching kits and  
5 conventional oral care products.

Conventional oral care products are products, such as tooth-pastes, gels, mouth washes or denture cleaning agents usually primarily target dental caries, plaque and/or tartar. However,  
10 oral care products also targeting tooth stains (often called tooth whiteners) have been available on the consumer market for some years. However, these products use another bleaching/whitening principle.

15 The above mentioned types of products are described further in the above section "Background of the Invention".

The oral composition or oral care product of the invention may comprise at least one other enzyme activity, which includes  
20 the activity of a protease, and/or mutanase and/or dextranase and/or lipase and/or amylase and/or anti-microbial polypeptides or enzymes.

In a preferred embodiment of the invention the oral composition or oral care product comprise an oxidoreductase and a  
25 dextranase and/or a mutanase.

An oral composition of the invention may advantageously be used for in conventional oral care products having any  
30 suitable physical form (i.e. powder, paste, gel, liquid, ointment, tablet etc.).

In an embodiments the oral composition or oral care products are one-compartment products in air-free packages containing  
35 an oxidoreductase enzyme or enzyme system, especially a laccase, a redox mediator agent (e.g. ABTS or PPT) and further ingredients normally used in such oral compositions or oral care products.

In another embodiment the oral care compositions or oral care products are one-compartment products comprising an oxidase and the corresponding substrate with a limited amount of water or a none-aqueous dentifrice.

Even though the presence of a redox mediator is advantageous, as it improves the action of the tooth bleaching, it is not compulsory.

Also contemplated according to the invention are two-compartment oral compositions and oral care products, where the oxidoreductase (e.g. laccase) and the redox mediator are mixed immediately before introduction into the oral cavity.

An "oral care product" of the invention is defined as a product which can be used for maintaining and/or improving oral hygiene in the mouth of humans and animals, and/or preventing or treating dental diseases.

Examples of such oral care products include toothpaste, dental cream, gel or tooth powder, odontic, mouth washes, denture cleaning agents, pre- or post brushing rinse formulations, chewing gum, lozenges, and candy.

Toothpastes and tooth gels typically include abrasive polishing materials, foaming agents, flavouring agents, humectants, binders, thickeners, sweetening agents, and water.

Mouth washes, including plaque removing liquids, typically comprise a water/alcohol solution, flavour, humectant, sweetener, foaming agent, and colorant.

According to the invention said abrasive polishing material includes alumina and hydrates thereof, such as alpha alumina trihydrate, magnesium trisilicate, magnesium carbonate, sodium bicarbonate ("Baking soda"), kaolin, aluminosilicates, such as calcined aluminum silicate and aluminum silicate, calcium car-

bonate, zirconium silicate, and also powdered plastics, such as polyvinyl chloride, polyamides, polymethyl methacrylate, polystyrene, phenol-formaldehyde resins, melamine-formaldehyde resins, urea-formaldehyde resins, epoxy resins, powdered  
5 polyethylene, silica xerogels, hydrogels and aerogels and the like. Also suitable as abrasive agents are calcium pyrophosphate, water-insoluble alkali metaphosphates, dicalcium phosphate and/or its dihydrate, dicalcium orthophosphate, tricalcium phosphate, particulate hydroxyapatite and the like.  
10 It is also possible to employ mixtures of these substances.

Dependent on the oral care product the abrasive product may be present in from 0 to 70% by weight, preferably from 1% to 70%. For toothpastes the abrasive material content typically lies  
15 in the range from 10% to 70% by weight of the final toothpaste product.

Humectants are employed to prevent loss of water from e.g. toothpastes. Suitable humectants for use in oral care products  
20 according to the invention include the following compounds and mixtures thereof: glycerol, polyol, sorbitol, polyethylene glycols (PEG), propylene glycol, 1,3-propanediol, 1,4-butanediol, hydrogenated partially hydrolysed polysaccharides and the like. Humectants are in general present from 0% to 80%,  
25 preferably 5 to 70% by weight in toothpaste.

Silica, starch, tragacanth gum, xanthan gum, extracts of Irish moss, alginates, pectin, cellulose derivatives, such as hydroxyethyl cellulose, sodium carboxymethyl cellulose and hydroxypropyl cellulose, polyacrylic acid and its salts, polyvinylpyrrolidone, can be mentioned as examples of suitable thickeners and binders, which help stabilizing the dentifrice product. Thickeners may be present in toothpaste creams and gels in an amount of from 0.1 to 20% by weight, and binders to  
30 the extent from 0.01 to 10% by weight of the final product.

As foaming agent soap, anionic, cationic, non-ionic, amphoteric and/or zwitterionic surfactants can be used. These

may be present at levels from 0% to 15%, preferably from 0.1 to 13%, more preferably from 0.25 to 10% by weight of the final product.

- 5 Surfactants are only suitable to the extent that they do not exert an inactivation effect on the present enzymes. Surfactants include fatty alcohol sulphates, salts of sulphonated mono-glycerides or fatty acids having 10 to 20 carbon atoms, fatty acid-albumen condensation products, salts of fatty acids  
10 amides and taurines and/or salts of fatty acid esters of isethionic acid.

Suitable sweeteners include saccharin and/or other appropriate sweeteners.

15

Flavours, such as spearmint and peppermint, are usually present in low amounts, such as from 0.01% to about 5% by weight, especially from 0.1% to 5%.

- 20 Water is usually added in an amount giving e.g. toothpaste a flowable form, i.e. between 40% to 70% by weight of the final product.

- 25 Further water-soluble anti-bacterial agents, such as chlorhexidine digluconate, hexetidine, alexidine, quaternary ammonium anti-bacterial compounds and water-soluble sources of certain metal ions such as zinc, copper, silver and stannous (e.g. zinc, copper and stannous chloride, and silver nitrate) may also be included.

30

- Also contemplated according to the invention is the addition of anti-calculus agents, anti-plaque agents, compounds which can be used as fluoride source, dyes/colorants, preservatives, vitamins, pH-adjusting agents, anti-caries agents, desensitizing agents etc.  
35

A toothpaste produced from an oral composition of the invention (in weight % of the final toothpaste composition)

may e.g. comprise the following ingredients:

	Abrasive material	10 to 70%
	Humectant	0 to 80%
5	Thickener	0.1 to 20%
	Binder	0.01 to 10%
	Sweetener	0.1% to 5%
	Foaming agent	0 to 15%
	Bleaching enzyme	0.0001% to 20%
10	Other enzymes	0 to 20%
	Peroxide	0 to 1%

#### Mouth washes

A mouth wash produced from an oral care composition of the invention (in weight % of the final mouth wash composition) may typically comprise the following ingredients:

	0-20%	Humectant
	0-2%	Surfactant
	0-5%	Enzymes
20	0-20%	Ethanol
	0-2%	Other ingredients (e.g. flavour, sweetener active ingredients such as florides).
	0-70%	Water

The mouth wash composition may be buffered with an appropriate buffer e.g. sodium citrate or phosphate in the pH-range 6-7.5.

The mouth wash may be in none-diluted form (i.e. must be diluted before use).

#### Method of Manufacture

The oral care composition and products of the present invention can be made using methods which are common in the oral product area.

Finally the invention relates to a method for using an oral care product of the invention, wherein

- a) the oral care product is introduced into the mouth,
- b) contacted with the teeth and/or gums for a period of time,
- c) removed from the mouth, and
- d) optionally rinsed with a liquid.

5

If the oral care product to be used is in solid to flowable form a tooth brush or the like may advantageously be used for contacting the oral care product with the teeth and/or gums. In the case of a liquid oral care product the contact may take place by rinsing the mouth.

10

The time period of contact in step b) is optional. However, contacting the oral care product with the teeth and/or gums for between about 30 seconds to 15 minutes will normally be sufficient for obtaining the desired result.

15

After use, the oral care product may be removed from the mouth in any suitable way, e.g. by spitting it out. Optionally the mouth may be rinsed with a liquid, such as tap water.

20

## METHODS AND MATERIALS

### Materials:

#### 25 Enzyme:

Laccase from *Myceliophthora thermophila*, (available from Novo Nordisk A/S).

Glucose oxidase from *Aspergillus niger* (available from Novo Nordisk A/S).

30

L-amino acid oxidase from *Trichoderma harzianum* (available from Novo Nordisk A/S).

#### 35 Teeth:

Alike coloured teeth for the bleaching tests are selected by colour determination on the Minolta CR-221 Chroma Meter. The teeth are stored in water under refrigeration until use.

Solutions:

Staining broth A: (extrinsic stains)

- 5 Clarified saliva\* 10 ml  
Tea\*\* 10 ml  
Coffee\*\*\* 10 ml  
Milk 10 ml

\* Clarified saliva is prepared by the following steps:

- 10 - chewing parafilm and collecting saliva,  
- centrifuging it and collecting the supernatant

\*\* Pour 50 ml of boiled water on to one Lipton™ Yellow label tea bag and leave it for 10 minutes.

\*\*\* Regular coffee

15

Staining broth B: (extrinsic stains)

100 ml autoclaved TSB (Tryptic Soy broth)

0.35 g instant coffee powder (Nescafe™, Classic)

0.25 g gastric mucin

20 Tea extract \*

3.25 ml TSB broth with 24 hours aerobically grown *Micrococcus luteus* DSM 20030T culture (from Nutrient agar plates grown for 2 days at 37°C).

- 25 \* 40 ml boiling water is poured to tea from one bag of Lipton™ Yellow label tea bag and are allowed to stand for 2 minutes. 15 ml of the supernatant is filtrated with a sterilized membrane, 0.45 µ.

Straining broth C: (intrinsic stains)

- 30 2.5% BSA in 0.1 M phosphate buffer, pH 7. 10 ml  
0.1 M Phosphate buffer, pH 7. 2.5 ml  
0.2% chlorhexidine digluconate 25 ml  
30% ribose 12.5 ml

Incubate at 37°C

35

Equipment:

Chroma Meter 210 (Minolta)

**Methods:**Preparation of hydroxyapatite tablets

Hydroxyapatite tablets are prepared by compressing 250 mg of hydroxyapatite in a tablet die at about 5,900 kg (13,000 lbs) of pressure for 5 minutes. The tablets are then sintered at 600°C for 4 hours and finally hydrated with sterile de-ionised water.

Sterilisation of hydroxyapatite tablets

HA tablets are sterilised at 180°C for two hours, hydrated with the sterilised de-ionised water and placed in a lid of Nunc tube (10 ml volume).

Staining of bovine teeth:

Bovine teeth are stained by immersing in staining broth A at 37°C until teeth surface become brown.

HA tablets stained with pellicle

HA tablets are immersed in 2 ml of the staining broth B placed in a Nunc tube, rotated with approximately 30° angle for 5 seconds with an interval of 10 seconds at 37°C. The staining broth is replaced every 24 hours. The HA tablets are incubated for 3 days.

Assessment of the teeth colour

Quantitative teeth colour assessment.

The colour of teeth and hydroxyapatite tablets are measured before and after the enzyme treatment on a Chroma Meter 210. The Chroma Meter has a 3 mm diameter circular aperture for measuring optical properties of small areas. The meter provides values of the optical parameters  $L^*$ ,  $a^*$  and  $b^*$  in the CIE system of colour measurement.  $\Delta L^*$  is the difference of  $L^*$  before and after a treatment which relates to the overall lightness or darkness change. Total colour difference is given by  $\Delta E^*$  calculated with an equation below. A high positive value of  $\Delta L^*$ , indicating lightness.



$$\Delta E^* = \sqrt{\Delta L^{*2} + \Delta a^{*2} + \Delta b^{*2}}$$

The colour of teeth should be measured after significant incubation of teeth in de-ionised water ( at least 15 minutes at room temperature ). The surface of teeth are wiped lightly with paper towel and the colour is measured.

L\*: "0"=black and "100"=white

#### Determination of Laccase Activity (LACU)

Laccase activity is determined from the oxidation of syringaldazin under aerobic conditions. The violet colour produced is photometered at 530 nm. The analytical conditions are 19  $\mu$ M syringaldazin, 23.2 mM acetate buffer, pH 5.5, 30°C, 1 minute reaction time.

1 laccase unit (LACU) is the amount of enzyme that catalyses the conversion of 1.0  $\mu$ mole syringaldazin per minute under these conditions.

#### Determination of peroxidase activity units (POXU)

Peroxidase activity is measured in POXU/ml. (1 POXU (peroxidase unit) is defined as the amount of enzyme that catalyses the conversion of 1  $\mu$ mole  $H_2O_2$  per minute in a system where 2,2'-azinobis[3-ethylbenzothiazoline-6-sulfonate] is oxidised in the presence of 1 mM  $H_2O_2$ , pH 7.0, at a temperature of 40°C.)

#### Determination of glucose oxidase activity (GODU)

1 GODU is defined as the amount of enzyme which, under standard conditions, catalyses the formation of 1 micromole of  $H_2O_2$  per minute. The analytic method AF266 is available upon request from Novo Nordisk A/S).

#### **EXAMPLES**

Example 1

### Bleaching of extrinsic teeth stains with glucose oxidase (GOX)

Bovine teeth were immersed in staining broth A comprising tea, coffee, milk, and saliva at 37°C until the teeth surface became brown as described in the Materials and Methods. section and treated with solution A and B, and Buffer as described below.

	Solution A Enzyme	Solution B Glucose	Buffer
1	none	none (Buffer)	50 mM Britton- Robinson pH 6.5
2	Glucose oxidase 50 GODU/ml	500 mM glucose	50 mM Britton- Robinson pH 6.0

Both samples were prepared with Britton-Robinson buffer (pH 6.5)  
Glucose solution was prepared one day before and kept at 4°C.  
The glucose oxidase was derived from *Aspergillus niger*

Each solution was pre-incubated at 37°C for 15 minutes prior to experiments

Two teeth were placed in a tube with 5.0 ml of solution A and 5.0 ml of solution B, respectively and incubation for 60 minutes at 37°C. A small magnetic stirrer was used for stirring the solutions. Both samples with buffer and oxidase + glucose, respectively, were treated this way. The treatment was repeated 3 times.

After incubation, the teeth were gently rinsed and the colour was measured as described in the Materials and Method section.

The result of the test is displayed in Table 1

Table 1

	$\Delta L^*$	$\Delta E^*$
Buffer	-0.15	3.13
25 GODU/ml glucose oxidase + 250 mM glucose	0.74	1.80

As can be seen from the Table 1 glucose oxidase is capable of

bleaching teeth with extrinsic stains.

### Example 2

#### 5 Bleaching of extrinsic stained hydroxy apatite tablets with L-amino acid oxidase

Hydroxyapatite (HA) tablets were extrinsically stained according to Stookey's method (Stookey G.K. et al. "In vitro Removal of Stains with Dentifrices", J. Dent. Res. 61, p. 10 1236 (1982)).

HA tablets (mimicking the properties of human enamel) sterilised as described in the Material and Methods section were immersed in staining broth B consisting of tea, coffee, 15 mucin and *Micrococcus luteus* for 5 seconds with 10 seconds intervals at 37°C as described above in the Materials and Method section. The HA tablets were incubated for 3 days. After staining the HA tablets were air dried and washed with de-ionised water

20 Chromatographically pure L-amino acid oxidase from *Trichoderma harzianum* free from catalase activity was incubated with a L-Arginine solution prepared in 0.1 M sodium phosphate buffer pH 8.5.

25 Two stained HA tablets were treated with said L-amino acid oxidase solution by incubating in a total volume of 0.5 ml at 37°C for 60 minutes with 90 reciprocal shaking/minutes.

30 The optical parameters L\*, a\* and b\* were measured on a Chroma Meter as described in the Materials and Methods section.

35 The result of the test is displayed in Table 2

Table 2

	$\Delta L^*$	$\Delta E^*$
Buffer	-0.17	0.30
6 Units/ml L-amino acid oxidase + 100 mM L-Arginine	1.04	1.07

As can be seen from the Table 2 L-amino acid oxidase is capable of bleaching teeth with extrinsic stains.

### 5 Example 3

#### Bleaching of extrinsic stained hydroxy apatite tablets with glucose oxidase

Hydroxyapatite (HA) tablets were stained according to Stookey's method (Stookey G.K. et al. " In vitro Removal of Stains with Dentifrices ", J. Dent. Res. 61, p. 1236 (1982)) and treated with chromatographically pure glucose oxidase from *Aspergillus niger*, free from catalase activity, incubated with a glucose solution prepared in 20 mM Britton-Robinson buffer, pH 6.0.

One stained HA tablet was treated with said glucose oxidase solution by incubating in a total volume of 10 ml at 39°C for 60 minutes with 90 reciprocal shaking/minutes.

The optical parameters L\*, a\* and b\* were measured on a Chroma Meter as described in the Materials and Methods section.

The result of the test is displayed in Table 3

Table 3

	$\Delta L^*$	$\Delta E^*$
Buffer	2.18	2.97
500mM glucose	1.80	1.84
10 GODU/ml glucose oxidase + 100 mM glucose	3.50	3.73
10 GODU/ml glucose oxidase + 500 mM glucose	3.62	3.62
50 GODU/ml glucose oxidase + 500 mM glucose	3.90	4.83
100 GODU/ml glucose oxidase + 500 mM glucose	5.05	5.63

As can be seen from the Table 3 glucose oxidase is capable of

bleaching teeth with extrinsic stains.

#### Example 4

#### Bleaching of extrinsic and intrinsic teeth stains with laccase

- 5 Bovine teeth were immersed in the staining broth (extrinsic stains) overnight at 37°C, transferred to staining broth c (intrinsic stains) and incubated at 37°C until the tooth surface became brown.

	Solution A Enzyme	Solution B mediator
1	none (Buffer)	none (Buffer)
2	0.6 LACU/ml Laccase	PTT 1000µM

- 10 Samples were all prepared with 50 mM Britton-Robinson buffer pH 6.5  
The laccase was derived from *Myceliophthora thermophila*

Each of solution A and B were pre-incubated at 37°C for 15 minutes prior to experiments

- 15 Two teeth were placed in a tube with 5.0 ml of solution A and 5.0 ml of solution B, respectively, and incubation for 60 minutes at 37°C. A small magnetic stirrer was used for stirring the samples. Both samples with buffer and Laccase +  
20 PPT, respectively, were treated this way. The treatment was repeated 3 times for each sample.

- After incubation, the teeth were gently rinsed and the colour was measured as described in the Materials and Method  
25 section.

The result of the test is displayed in Table 4

Table 4

	$\Delta L^*$	$\Delta E^*$
Buffer	0.01	1.53
0.3 LACU/ml laccase + 500 µM PPT	2.67	2.78

- 30 As can be seen from the Table 4 laccase is capable of bleaching teeth with intrinsic stains.

As will be apparent to those skilled in the art in the light of the foregoing disclosure, many alterations and modifications are possible in the practice of this invention without departing from the spirit or scope thereof. Accordingly, the scope of the invention is to be construed in accordance with the substance defined by the following claims.

**PATENT CLAIMS**

1. An oral composition for bleaching teeth comprising at least one oxidoreductase.  
5
2. The oral composition according to claim 1, comprising at least one enzyme capable of using molecular oxygen (O<sub>2</sub>) or a peroxide as acceptor.
- 10 3. The oral composition according to claims 1 and 2, wherein the enzyme is laccase or related enzymes.
- 15 4. The oral composition according to claim 3, wherein the laccase is derived from *Polyporus* sp., in particular *Polyporus pinsitus*, or *Myceliophthora*, in particular *M. thermophila*, or *Scytalidium*, in particular *S. thermophilum*.
- 20 5. The oral composition according to claim 3, wherein the laccase related enzyme is a catechol oxidase or a bilirubin oxidase.
6. The oral composition according to claims 1 and 2, wherein the enzyme is an oxidase.
- 25 7. The oral composition according to claim 6, wherein the oxidase is glucose oxidase
- 30 8. The oral composition according to claim 6, wherein the oxidase is selected from the group of oxidases including hexose oxidase, L-amino-acid oxidase, xylitol oxidase, galactose oxidase, pyranose oxidase, and alcohol oxidase.
- 35 9. The oral composition according to claims 1 and 2, wherein the enzyme is a peroxidase or a haloperoxidase.
10. The oral composition according to any of claims 1 to 9, wherein the oral composition comprises a laccase or a related enzymes and/or an oxidase and/or a peroxidase.

11. The oral composition according to any of claims 1 to 10,  
wherein the oxidoreductase is present in a concentration of  
0.0001% to 20%, preferably 0.001 to 10% especially 0.01% to 5%  
5 calculated on the basis of the oral composition.
12. The oral care composition according to any of claims 1 to  
10, comprising a from 0% to 0.5% peroxide, such as 0.1% to  
0.3%, preferably 0% to 1% calculated of the basis of the final  
10 oral composition.
13. The oral composition according to any of claims 1 to 12,  
comprising a mediator, such as ABTS or PPT.
14. The oral care composition according to any of claims 1 to  
13, wherein the bleaching enzyme can act on substrates which  
are not cariogenic.
15. The oral care composition according to any of claims 1 to  
14, wherein the bleaching enzyme(s) is substantially active in  
the pH range from between 5.0 to 9.0, preferably between pH  
20 6.0 to 8.5, especially between pH 6.4 to 7.5.
16. An oral care product, comprising an oral composition for  
25 bleaching teeth according to any of claims 1 to 15.
17. An oral care product according to claim 16, being a den-  
tifrifice, such as a toothpaste, tooth powder, a mouth wash, a  
pre-rinse or a denture cleaning agent.
- 30 18. The oral care product according to claims 16 and 17,  
comprising an anti-plaque compound or composition and/or an  
anti-tartar compound or composition.
- 35 19. The oral care product according to any of claims 16 to 18,  
comprising at least one other enzyme activity selected from  
the group including a protease and/or amylase and/or  
dextranase and/or mutanase and/or lipase and/or an anti-



microbial activity, or a mixture thereof.

20. The oral care product according to any of claims 16 to 19,  
comprising at least one oxidoreductase and a dextranase and/or  
5 a mutanase.

21. The oral care product according to any of claims 16 to 20,  
comprising at least a laccase and a dextranase and/or a muta-  
nase.

10 22. The oral care product according to any of claims 16 to 21,  
wherein the oxidoreductase is present in a concentration of  
0.0001% to 20%, preferably 0.001 to 10% especially 0.01% to 5%  
calculated on the basis of the final oral care product.

15 23. The oral care product according to any of claims 16 to 22,  
comprising from 0% to 1%, such as between about 0.1% to 0.5%,  
preferably 0% to 3% peroxide, especially 0% to 6% calculated  
of the basis of the weight of the final oral care product.

20 24. A method for bleaching discoloured or stained teeth, com-  
prising  
a) subjecting the teeth to an oral composition according to  
any of claims 1 to 15 or an oral care product according to any  
25 of claims 16 to 23.

25. A method for using an oral composition according to any of  
claims 1 to 15 or an oral care products according to any of  
claims 16 to 23 in the oral cavity, wherein  
30 a) the oral care product is introduced into the mouth,  
b) contacted to the teeth and/or gums for a period of time,  
c) removed from the mouth, and  
d) optionally rinsed with a liquid.

35 26. The method according to claim 25, wherein a tooth brush is  
used for contacting the oral care product with the teeth.

27. Use of oxidoreductases for bleaching teeth.

28. The use according to to claim 27, wherein the oxidoreductase is selected from the group including laccases and related enzymes, and oxidases, such as glucose oxidase, hexose oxidase, L-amino-acid oxidase xylitol oxidase, galactose oxidase, pyranose oxidase, and alcohol oxidase, and peroxidases, such as a haloperoxidase.
29. The use according to claim 28 for bleaching non-vital teeth.
30. The use according to claim 28 for bleaching vital teeth.
31. The use according t claim 30 in over-the-counter tooth product.
32. The use according to claim 31, wherein the over-the-counter-tooth product is an oral composition according to any of claims 1 to 15 or an oral care products according to any of claims 16 to 23.
33. The use according to claim 31, wherein the oral care product is dentifrice, such as a toothpaste, mouth rinse or pre-rinse or denture cleaning agents.
34. The use according to claim 31, in a home-applied tooth bleaching composition or tooth bleaching product
35. The use according to claim 30, in "in-office"-tooth bleaching compositions or tooth bleaching product.
36. The use according to any of claims 27 to 35, wherein the oxidoreductase is an enzyme according to claims 2 to 15, or a mixture thereof.

1  
INTERNATIONAL SEARCH REPORT

International application No.  
PCT/DK 96/00350

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC6: A61K 7/28

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC6: A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, CA, BIOSIS

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4150113 A (HENDRIK HOOGENDOORN ET AL), 17 April 1979 (17.04.79), abstract; claims 1 and 7 --	1-36
X	EP 0133736 A2 (LACLEDE PROFESSIONAL PRODUCTS, INC.), 6 March 1985 (06.03.85), abstract; claims 1-8 --	1-36
X	DE 2027019 A (TELEC S.A.), 17 December 1970 (17.12.70), claims 1-4 --	1-36

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

\* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

21 November 1996

Date of mailing of the international search report

22 -11- 1996

Name and mailing address of the ISA/  
Swedish Patent Office  
Box 5055, S-102 42 STOCKHOLM  
Facsimile No. +46 8 666 02 86

Authorized officer

Dagmar Järvman  
Telephone No. +46 8 782 25 00

2  
INTERNATIONAL SEARCH REPORT

International application No.  
PCT/DK 96/00350

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>Dialog Information Services, File 350, Dialog accession no. 002242548, WPI accession no. 79-41744B/22, Anonymous: "Tooth cleaning agent contg. mutanase and glucose oxidase - with good plaque decomposing and spot removing action; PASTE", RD,A,181075, 790510, 7922 (Basic)</p> <p style="text-align: center;">--</p>	1-36
X	<p>US 4269822 A (MICHAEL A. PELLICO ET AL), 26 May 1981 (26.05.81), abstract; claims 1 and 3</p> <p style="text-align: center;">-- -----</p>	1-36

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK96/00350

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. Oral composition comprising oxidoreductase according to claims 1,2 and 24-27
2. Oral composition comprising laccase according to claims 3-5 and 24-27

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

28/10/96

International application No.  
PCT/DK 96/00350

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A- 4150113	17/04/79	AT-A,B- 306243 BE-A- 751390 CA-A- 926777 CH-A- 531885 DE-A- 2027019 FR-A- 2045815 GB-A- 1309282 NL-B- 150332 NL-A- 6908379 SE-B- 366651 US-A- 4178362	15/02/73 03/12/70 22/05/73 31/12/72 17/12/70 05/03/71 07/03/73 16/08/76 07/12/70 06/05/74 11/12/79
EP-A2- 0133736	06/03/85	JP-C- 1740554 JP-B- 4025924 JP-A- 59231011 US-A- 4537764 US-A- 4564519	15/03/93 06/05/92 25/12/84 27/08/85 14/01/86
DE-A- 2027019	17/12/70	AT-A,B- 306243 BE-A- 751390 CA-A- 926777 CH-A- 531885 FR-A- 2045815 GB-A- 1309282 NL-B- 150332 NL-A- 6908379 SE-B- 366651 US-A- 4150113 US-A- 4178362	15/02/73 03/12/70 22/05/73 31/12/72 05/03/71 07/03/73 16/08/76 07/12/70 06/05/74 17/04/79 11/12/79
US-A- 4269822	26/05/81	NONE	